

Centrally condensed turbulent cores: Massive stars or fragmentation?

Clare L. Dobbs,^{*} Ian A. Bonnell and Paul C. Clark

School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, Fife, KY16 9SS

Accepted

ABSTRACT

We present numerical investigations into the formation of massive stars from turbulent cores of density gradient $\rho \propto r^{-1.5}$. The results of five hydrodynamical simulations are described, following the collapse of the core, fragmentation and the formation of small clusters of protostars. We generate two different initial turbulent velocity fields corresponding to power-law spectra $P \propto k^{-4}$ and $P \propto k^{-3.5}$, and apply two different initial core radii. Calculations are included for both completely isothermal collapse, and a non-isothermal equation of state above a critical density (10^{-14}gcm^{-3}). Our calculations reveal the preference of fragmentation over monolithic star formation in turbulent cores. Fragmentation was prevalent in all the isothermal cases. Although disc fragmentation was largely suppressed in the non-isothermal runs due to the small dynamic range between the initial density and the critical density, our results show that some fragmentation still persisted. This is inconsistent with previous suggestions that turbulent cores result in the formation of a single massive star. We conclude that turbulence cannot be measured as an isotropic pressure term.

Key words: stars: formation – turbulence – hydrodynamics

1 INTRODUCTION

There are several potential difficulties in forming high-mass stars. Firstly, the timescale of less than 10^6 years to assemble 10 to more than $100 M_{\odot}$ implies large accretion rates (Zinnecker et al. 1993; Behrend & Maeder 2001; Norberg & Maeder 2000). Secondly, their crowded location in the centre of young stellar clusters (Clarke et al. 2000; Hillenbrand & Hartmann 1998; Carpenter et al. 1997; Lada & Lada 2003) limits the final mass of any collapsing fragment: confining the Jeans radius to be less than the interstellar separation means that the resultant Jeans mass will be correspondingly small (Zinnecker, McCaughrean & Wilking 1993; Bonnell, Bate & Zinnecker 1998). Lastly, and most importantly, the radiation pressure from a high-mass star is sufficient to reverse the infall of gas that contains typical dust properties (Wolfire & Cassinelli 1987; Beech & Mitalas 1994). There are a number of ways this last problem can be circumvented. The radiation pressure could be overwhelmed by ultra-high accretion rates (McKee & Tan 2003). Alternatively, accretion could occur preferentially through an equatorial disc, while the central object is rapidly rotating and thus emitting most of its radiation towards the poles (Yorke & Sonnhalter 2002). A third solution is that massive

stars form due to stellar mergers in the ultra-dense core of a cluster (Bonnell et al. 1998; Bonnell & Bate 2002).

The McKee & Tan (2003, 2002) model depends on high accretion rates that would be expected to occur in a dense gas core in the centre of a cluster. This core is envisioned to be supported by turbulence, as thermal pressure is inadequate, and adopts a steep density profile in order to prevent fragmentation. Even neglecting how such a core could arise in the centre of a stellar cluster, one potential difficulty is that turbulent clouds are known for their tendency to fragment and form a stellar cluster. It is this possibility that we address here.

Fragmentation during the initial stages of star formation has been illustrated through numerical studies of collapsing turbulent molecular clouds. Bate, Bonnell & Bromm (2003) describe the collapse of a uniform $50 M_{\odot}$ turbulent cloud of diameter 0.375pc , which forms 3 cores containing 23+ protostars and 27+ brown dwarfs. Fragmentation has also been demonstrated in isolated, rotating $1 M_{\odot}$ cloud cores (Burkert, Bate & Bodenheimer 1997), where low multiple systems are produced despite an initial $\rho \propto r^{-1}$ density distribution.

The formation of stellar clusters from turbulent cloud cores (Bonnell, Bate & Vine 2003) has shown how the fragmentation of the turbulently induced filamentary structure produces hundreds of stars. These stars fall into local poten-

^{*} E-mail: cld2@st-and.ac.uk

tial wells forming small sub-clusters which eventually merge to form a large stellar cluster. The massive stars form in the centre of the sub-clusters due to competitive accretion, naturally explaining why massive stars are found in the cores of dense stellar clusters (Bonnell, Vine & Bate 2004). The stellar densities in these cores can be very high, approaching that for stellar mergers to occur (Bonnell & Bate 2002).

Our calculations adopt a turbulent core with a density profile and velocity dispersion comparable to McKee & Tan (2002). The purpose of this paper is to investigate whether the steep density gradient assumed by McKee and Tan is sufficient to prevent fragmentation and consequently provide a suitable approach to massive star formation.

2 COMPUTATIONAL METHOD

2.1 SPH code

We use the 3D smoothed particle hydrodynamics (SPH) code based on the version by Benz (Benz 1990). The smoothing lengths between particles are allowed to vary, but the typical number of neighbours for each particle is $N_{\text{neigh}} \sim 50$. Artificial viscosity is included with the standard parameters $\alpha = 1$ and $\beta = 2$. Gravitational forces are calculated using a binary tree. The code uses 10^6 particles (so for a $30M_{\odot}$ core (Section 2.3) the minimum particle mass is $3 \times 10^{-5}M_{\odot}$) and simulations were run for approximately half a free-fall time. All computations were performed using the United Kingdom's Astrophysical Fluids Facility (UKAFF), a 128 CPU SGI Origin 3000 supercomputer.

2.2 Sink particles

Simulations in SPH become very computationally expensive as the density increases during the collapse of a core. The insertion of sink particles (Bate, Bonnell & Price 1995) at a certain density is widely used to extend calculations. To resolve the local Jeans mass requires $\sim 2N_{\text{neigh}}$ particles (Bate & Burkert 1997), i.e. $3 \times 10^{-3}M_{\odot}$ in these simulations. This is then the minimum resolvable mass that can be considered unstable for collapse. The minimum density for insertion of a sink particle can then be determined from Jean's equation, inserting $3 \times 10^{-3}M_{\odot}$ as the Jean's mass:

$$M_J = \left(\frac{5R_g T}{2\mu G} \right)^{3/2} \left(\frac{4}{3}\pi\rho \right)^{-1/2} = 3 \times 10^{-3}M_{\odot} \quad (1)$$

Taking the temperature T to be $20K$ and $\mu = 2.46$, this rearranges to give $\rho_{\text{acc}} = 1.06 \times 10^{-13}\text{gcm}^{-3}$. The corresponding radius for accretion is $r_{\text{acc}} \sim 20\text{AU}$.

2.3 Initial conditions

McKee & Tan (hereafter MT(2003)) assume a centrally condensed core, following a density profile of $\rho \propto r^{-k_{\rho}}$. We take $k_{\rho} = 1.5$, the fiducial value of MT(2003). This is shallower than the singular isothermal sphere model ($k_{\rho} = 2$, (Shu 1977)) but comparable with observations of cloud cores (Myers & Fuller (1992) give $k_{\rho} = 1.6$, Ward-Thompson et al. (1994) $k_{\rho} = 1.2$).

The first computations took the core radius R to be

0.06pc with a mass of $30M_{\odot}$. With a resolution of 10^6 particles, the initial density profile $\rho \propto r^{-3/2}$ rendered a central density of $\sim 3 \times 10^{-15}\text{gcm}^{-3}$, whilst the average density was $\rho_{\text{av}} \sim 2.5 \times 10^{-18}\text{gcm}^{-3}$. The results are given in terms of the free fall time, $t_{\text{ff}} = (3\pi/32G\rho_{\text{av}})^{-1/2} \sim 4.5 \times 10^4$ years. MT(2003) use similar initial conditions, but take a mass of $60M_{\odot}$ to form a $30M_{\odot}$ star with 50% efficiency. We also performed computations with a core radius 0.2pc, since this gave similar dimensions to Bate et al. (2003). The average density was then $3 \times 10^{-19}\text{gcm}^{-3}$ and the central density $2 \times 10^{-16}\text{gcm}^{-3}$. The free fall time for these results was 2.7×10^5 years. In all calculations, the initial temperature of the core was 20K, comparable to the cold gas of which the first massive stars form.

As in equation (1), the Jeans mass is

$$M_J(\rho(r)) = \left(\frac{5R_g T}{2\mu G} \right)^{3/2} \left(\frac{4}{3}\pi\rho \right)^{-1/2} \quad (2)$$

where μ the mean molecular weight is taken as 2.46. The number of Jeans masses contained in a core is then

$$N_J(R) = \int_0^R \frac{\rho(r)4\pi r^2}{M_J(\rho(r))} dr. \quad (3)$$

This gives 69 Jean's masses when $R = 0.06\text{pc}$ and 11.5 when $R = 0.2\text{pc}$.

2.3.1 Turbulence

MT(2003) incorporate turbulence by means of an effective turbulent pressure term whereby $P \propto r^{-k_P}$. By assuming hydrostatic equilibrium, $k_P = 2(k_{\rho} - 1)$, and the resulting velocity dispersion relation is derived as

$$\sigma \propto c_s = \sqrt{\frac{P}{\rho}} \propto r^{(2-k_{\rho})/2} \quad (4)$$

where σ is the velocity dispersion, and c_s the effective isothermal sound speed. Inserting $k_{\rho} = 1.5$ then gives $\sigma \propto r^{0.25}$. This is shallower than observational results obtained for the similar Larson relation $\sigma \propto L^{\alpha}$ (L a typical length scale), which indicate that $0.25 < \alpha < 0.75$, e.g. $\alpha = 0.38$ (Larson 1981), $\alpha = 0.5$ (Myers 1983).

We simulate turbulence using a grid method outlined by Dubinski et al. (1995) and briefly described here. The initial velocity field is calculated from a random Gaussian distribution and follows a power spectrum

$$P(k) \equiv \langle |v_k|^2 \rangle \propto k^{-n}, \quad (5)$$

where k is the wavenumber. To obtain a divergence free velocity field, velocities are constructed from a vector potential $\mathbf{A}$, such that $\mathbf{v} = \nabla \times \mathbf{A}$. We sample components of $\mathbf{A}_{\mathbf{k}}$, the Fourier transform of $\mathbf{A}$, from a grid of co-ordinates (k_x, k_y, k_z) . In order to generate a Gaussian distribution satisfying (5), components of $\mathbf{A}_{\mathbf{k}}$ are selected as

$$\mathbf{A}_{\mathbf{k}}(k_x, k_y, k_z) = |\mathbf{k}|^{\frac{-n-2}{2}} (C_{k_x} e^{i\theta_{k_x}}, C_{k_y} e^{i\theta_{k_y}}, C_{k_z} e^{i\theta_{k_z}}) \quad (6)$$

The amplitudes C are sampled from a Rayleigh distribution and phases $0 < \theta < 2\pi$ from a random distribution (see Dubinski et al. (1995)). Taking the inverse Fourier transform of $\nabla \times \mathbf{A}_{\mathbf{k}}$ then gives

$$\mathbf{v} = \frac{1}{2\pi} \sum_{\mathbf{k}} i\mathbf{k} \times \mathbf{A}_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}} \quad (7)$$

Numerically the real x component of (7) becomes

$$v_x = \frac{1}{2\pi} \sum_{-k_{max}}^{k_{max}} |\mathbf{k}|^{\frac{-n-2}{2}} \left[k_z C_{k_y} \sin(\mathbf{k} \cdot \mathbf{r} + \phi_{k_y}) - k_y C_{k_z} \sin(\mathbf{k} \cdot \mathbf{r} + \phi_{k_z}) \right] \quad (8)$$

summed over k_x, k_y and k_z . Similar expressions provide v_y and v_z , for a particle at position (x, y, z) . The parameter k_{max} in equation (8) is often chosen to satisfy the resolution requirements of the model, i.e $k_{max} \sim (2\pi/L_{min})$ where L_{min} is the smallest length scale or minimum separation between particles. However, since our model is centrally condensed and the density varies considerably, we allow k_{max} to vary for each particle according to the local smoothing length. We take

$$(k_{max})_i = \text{INT} \left(\frac{2\pi}{h_i} \right) \quad (9)$$

h_i is the smoothing length of the i th particle. In this way, velocities in less dense (low resolution) regions, could be computed much faster than those in the dense centre of the core (high resolution), reducing the computational expense of the calculation.

The velocity field generated gives a dispersion - length scale dependence similar to the Larson relation. Calculations are included for $\sigma \propto r^{0.25}$, consistent with MT(2003), and $\sigma \propto r^{0.5}$ which corresponds better with observations. These dispersion laws are attained by generating power spectra with $n = 3.5$ and $n = 4$ respectively. However Myers & Gammie (1999) show that $n \sim 3.25$ provides a better fit for the $\sigma \propto r^{0.25}$ dispersion law.

The core is initially in virial equilibrium, such that

$$E_{grav} + 2E_{turb} + 2E_{therm} = 0. \quad (10)$$

The ratios of turbulent to gravitational energy were

$$\frac{E_{turb}}{E_{grav}} \sim 0.4 \quad \text{for } R = 0.06pc, \quad (11)$$

$$\frac{E_{turb}}{E_{grav}} \sim 0.25 \quad \text{for } R = 0.2pc. \quad (12)$$

The turbulent energy is then allowed to dissipate over the dynamical timescale of the core (Mac Low et al. 1997). We do not include any driving, which if included, should increase the susceptibility to fragmentation.

2.3.2 Equation of State

The thermal behaviour of the cloud core was assigned as either:

- (i) Completely isothermal.
- (ii) Non-isothermal above a critical density, allowing the gas to heat up as the core collapses. In this case the equation of state exponent varies as

$$\gamma = \begin{cases} 1 & \text{if } \rho < 10^{-14} \text{gcm}^{-3} \\ 1.67 & \text{if } \rho \geq 10^{-14} \text{gcm}^{-3} \end{cases} \quad (13)$$

- (iii) Non-isothermal above the critical density with a second isothermal collapse phase for $\rho > 10^{-12} \text{gcm}^{-3}$.

Although the core is initially globally in virial equilibrium, the inner regions are actually over-supported due to the isothermal nature of the gas.

3 RESULTS

A summary of the calculations undertaken, their parameters and the overall results are shown in Table 1. The parameter α describes the initial velocity dispersion (where $\sigma \propto r^\alpha$).

In all simulations, turbulence initially supports the core in virial equilibrium. As the simulation progresses, turbulence decays through shocks in the core, dissipating kinetic energy. The core then begins to undergo collapse, with density increasing until the formation of individual protostars. The morphology of the cloud is similar in each simulation - the core becomes elongated, generating significant structure which forms the basis for subsequent fragmentation (e.g. Figure 1).

3.1 Isothermal results

Models 1, 2 and 3 used an isothermal equation of state throughout the simulation. Models 1 & 3 apply a $\sigma \propto L^{0.25}$ dispersion law whereas Model 2 uses $\sigma \propto L^{0.5}$ although the initial velocity fields are determined by the same series of random numbers in all calculations. Model 3 differs by taking a 0.2pc radius cloud.

All 3 simulations produced significant numbers of protostars. Table 1 includes the total number of protostars formed, although results cannot be directly compared as some formation was still ongoing.

3.1.1 Comparison between dispersion laws

Figure 1 displays different stages during Models 1 & 2, which apply different power spectra. The main difference between Models 1 & 2 is the timescale for protostar formation and fragmentation. Collapse and protostar formation occur later in Model 1 compared to Model 2, although their overall evolution is similar. This is due to the shallower power law $P \propto k^{-3.5}$ (compared to $P \propto k^{-4}$ for Model 2 when $\alpha = 0.5$) which supplies more kinetic energy over smaller length scales. This provides greater support over small scales and generates more structure. Consequently density plots for Model 1 show more structure than those for Model 2, and the protostars are less widely distributed (Figure 1).

The left side of Figure 1 illustrates the evolution for the $\alpha = 0.25$ dispersion law, which best represents MT(2003). A protostar forms in the central part of the core after $0.19t_{ff}$, surrounded by a disc of gas. The core fragments after $0.25t_{ff}$, when 2 further protostars form from the disc material. A fourth protostar is also created via fragmentation of an elongated filament extending from the initial protostar (Figure 1 ii). No further protostars form for $\sim 0.15t_{ff}$, as the core is largely supported by internal rotation. However the internal dynamics are still chaotic and further collapse occurs as support is lost. Further fragmentation occurs through collapse along filaments near the core's centre (Figure 1 iii). This results in a second phase of star formation at around $0.47t_{ff}$, producing a total of 28 protostars after $0.54t_{ff}$.

Model 2 (Figure 1, right), where $\alpha = 0.5$, shows similar behaviour. Again an initial protostar is formed after $0.09t_{ff}$ accompanied by a disc which subsequently distorts and fragments, producing 7 protostars after $0.13t_{ff}$. Similarly there is a second phase of fragmentation through the collapse of




Figure 1. Column density plots (logarithmic scale, minimum and maximum densities of $4 \times 10^{-17} \text{gcm}^3$ and $4 \times 10^{-13} \text{gcm}^3$) showing evolution of central region of the core for $\sigma \propto L^{0.25}$ (left) and $\sigma \propto L^{0.5}$ (right). Length scale for each plot is $0.02 \text{pc} \times 0.02 \text{pc}$. The number of protostars in the plots are: Left - i)1, ii)4, iii)6, iv)13; Right - i)1, ii)7, iii)7, iv)14. Times are given in units of $t_{ff} = 4.5 \times 10^4$ years.

Model	α	R (pc)	Jean No.	Predominant time of formation (t_{ff})	No. of protostars	Total mass accreted ($M_\odot$)	Most massive protostar ($M_\odot$)
1	0.25	0.06	95	0.47	28	2.73	0.435
2	0.5	0.06	95	0.37	14	2.12	0.528
3	0.25	0.2	16	0.55	19	1.79	0.66
4	0.25	0.06	95	0.54	3 [†]	2.02	0.99
5	0.25	0.2	16	0.54	2	2.10	1.28

Table 1. Table showing results of all simulations, giving the total number of stars formed and the approximate time at which most formation occurred. Models 4 & 5 use a non-isothermal equation of state above a critical density (Section 2.2.2). However Model 4 also includes a second isothermal collapse phase. [†]Further fragmentation apparent but sink formation suppressed by equation of state.

elongated filaments, leading to a total of 14 protostars after $0.44t_{ff}$. Figure 2 plots the mass accreted and the number of protostars formed against fraction of a free fall time, for both dispersion laws. Models 1 and 2 display very similar profiles, offset by $\sim 0.1t_{ff}$. The number of protostars formed in each case shows 2 periods of star formation at $0.1/0.2t_{ff}$ and $0.38/0.48t_{ff}$ separated by a period of $0.26/0.15t_{ff}$.

3.1.2 Overall comparison of isothermal runs

The isothermal results all display a high level of fragmentation, demonstrating that turbulent pressure is anisotropic. Turbulence cannot be represented by an isotropic thermal pressure, instead distortion of the core leads to the formation of multiple protostars. The degree and timescale of fragmentation vary according to the turbulent power law and the size of the core. Whereas a steeper velocity-sizescale relation ($\alpha = 0.25$) advances protostellar formation, collapse is delayed for the $R = 0.2\text{pc}$ core in Model 3. The larger core has more thermal support so the ratio of thermal to gravitational energy is greater (equations (11),(12)). Protostar formation begins at $\sim 0.52t_{ff}$, so collapse of the core and subsequent fragmentation occurs at a higher fraction of the free fall time compared with Model 1 where $R = 0.06\text{pc}$. Correspondingly, density plots for Model 3 also show less structure, tending to retain a centrally condensed profile, at least over large length scales. There was, however, still sufficient structure over the centre to produce 19 protostars over $0.6t_{ff}$, so significant fragmentation still occurred.

3.1.3 Accretion rates

The main motivation for MT(2003) was to achieve a high accretion rate to overcome radiation pressure. Accretion rates can generally be estimated as the resulting protostellar mass divided by the free-fall time (M/t_{ff}). Thus MT(2003) construct their models to produce an early accretion rate of $10^{-4}M_\odot\text{yr}^{-1}$, which increases to $10^{-3}M_\odot\text{yr}^{-1}$. From Figure 2 we see the accretion rate is initially $\sim 10^{-4}M_\odot\text{yr}^{-1}$ increasing to $10^{-3}M_\odot\text{yr}^{-1}$ after $0.3t_{ff}$. These accretion rates agree with MT(2003), but are based on the total mass accreted onto many protostars (between ~ 5 -20). The accretion rate for an individual protostar is therefore significantly less. In contrast, the cluster accretion models for massive star formation (Bonnell et al. 2004) show that although the mean accretion rate (M/t_{ff}) is only $10^{-6}M_\odot\text{yr}^{-1}$, the actual accretion rate onto the growing massive star in the centre of a cluster is $10^{-4}M_\odot\text{yr}^{-1}$.




Figure 2. Plots for number of stars (top) and mass accreted (bottom) versus time for Model 1, $\alpha = 0.25$ (solid line) and Model 2, $\alpha = 0.5$ (dashed line).

3.2 Non isothermal equation of state

The large scale collapse of the non-isothermal models are very similar to the corresponding isothermal cases, since the equation of state changes only in the central denser regions. However, Table 1 shows that the number of protostars formed is dramatically reduced (Models 4 & 5). In Model 4, (0.06pc core), the equation of state changed before fragments could collapse to form sink particles. A second isothermal collapse phase subsequently produced 3 protostars. For Model 5 (0.2pc core), the core is initially less dense so 2 protostars were able to form without any second collapse phase.

To understand the difference between protostar formation in the isothermal and non-isothermal models, it is necessary to compare the protostar-forming region in each case. Figure 3 illustrates the contrast between Model 1 (isothermal) and Model 4 (non-isothermal) in the dense part of the core. Complex structures continue to exist on small scales in the isothermal case, including a disc and close companions formed through disc fragmentation. In comparison, Model 4 is dominated by 2 large volumes of dense thermally supported gas.

On scales comparable to the core radius however, density profiles show that considerable structure still exists in the non-isothermal core. In Figure 4 (Model 4), the maximum densities over the x and y directions are plotted (while the equation of state remains non-isothermal). Five peaks have been selected from the profiles which show a large density contrast with the surrounding material. These indicate independent, disconnected fragments. Figure 5 shows an accompanying column density plot, obtained at the same time frame, which illustrates the large scale structure of the core. The locations of the five peaks are indicated, showing where fragmentation is taking place. The 2 highest peaks in the densest part of the core correspond to the dense regions shown in Figure 3. These also subsequently form 3 protostars during the second isothermal collapse phase, as the larger region (of Figure 3) sub-fragments. With a longer period of time, and a less severe equation of state, we would expect all the dense fragments corresponding to these peaks to collapse independently and form protostars.

The distortion of the 0.06pc core suggests that monolithic formation is unlikely, despite the formation of very few protostars. Our analysis suggests fragmentation will occur producing at least 6 separate bodies (5 peaks identified from Figures 4 & 5, and isothermal sub-fragmentation of peak 2). This is again a consequence of anisotropic turbulence dominating homogeneous thermal support.

Conversely, the larger core (Model 5) remains predominantly centrally condensed. Thermal pressure supports the core over large scales, whilst the equation of state prevents any local fragmentation. With only 2 centrally located protostars, Model 5 would only be expected to produce a low multiple (2 or 3 protostars) system.

4 CONCLUSIONS

Turbulently supported clouds of typical density structure $\rho \propto r^{-1.5}$ have been hypothesised to be the progenitors of massive stars. Their density concentration could prevent fragmentation and their high accretion rates overwhelm the radiation pressure from the accreting massive star (McKee & Tan 2003, 2002). We have performed numerical simulations of this scenario and found that the initial centrally condensed density profile proved insufficient to prevent fragmentation. Turbulent support generates significant structure in the core which forms the basis for subsequent fragmentation. Turbulence cannot be assumed to act as the equivalent of an isotropic pressure. In addition, although the total mass accretion rates are comparable with MT(2003), individual protostellar accretion rates are significantly lower.

The fragmentation of the centrally condensed $30M_{\odot}$ cores forms ≈ 20 stars over the time period of our isother-




Figure 3. Logarithmic column density plots (as Figure 1, minimum and maximum densities of $4 \times 10^{-17} \text{gcm}^{-3}$ and $4 \times 10^{-13} \text{gcm}^{-3}$) when $\alpha = 0.25$, $R=0.06\text{pc}$ for isothermal (Model 1, top) and non-isothermal (Model 4, bottom) cases. The size of the plots are 0.02pc by 0.02pc and the time is $0.32t_{ff}$, $t_{ff} = 4.5 \times 10^4$ years.

mal simulations. The protostars form from a combination of the fragmentation of filamentary structure and the fragmentation of rotationally supported discs. Our non-isothermal runs suppress this disc fragmentation as the gas is assumed to heat on scales of the discs. Nevertheless, additional fragmentation is expected once a second collapse phase softens the equation of state. Competitive accretion in these clusters will determine the eventual stellar masses (Bonnell et al. 2001).

These simulations neglect the potential effects of any magnetic fields present in the core. We do note that this should not impede the fragmentation process as magnetic fields have been shown not to affect the structure generation in turbulent molecular clouds (Stone et al. 1998; Padoan et al. 2004). Even in the presence of magnetic fields, turbulence does not act as an isotropic support, and neither




Figure 4. Logarithmic plot of maximum density (over all points in the simulation) versus x (red) and versus y (green) for Model 4. Dotted lines show 5 dominant peaks. ρ measured in computational units where 1 unit = $7 \times 10^{-17} \text{ g cm}^{-3}$. The plot is taken at $t = 0.53t_{ff}$ ($t_{ff} = 4.5 \times 10^4$ years) when the equation of state is non-isothermal.




Figure 5. Column density plot for Model 4, with peaks identified in Figure 4 indicated. Only points where $\log(\rho) > -2$ have been selected to emphasise the structure. The size scale of this plot is 0.1 pc by 0.1 pc with minimum and maximum densities of $7 \times 10^{-19} \text{ g cm}^{-3}$ and $4 \times 10^{-13} \text{ g cm}^{-3}$. Again the plot corresponds to a time of $t = 0.53t_{ff}$.

turbulence nor magnetic fields should be modelled as an isotropic pressure term.

REFERENCES

- Bate M. R., Bonnell I. A., Bromm V., 2003, MNRAS, 339, 577
- Bate M. R., Bonnell I. A., Price N. M., 1995, MNRAS, 277, 362
- Bate M. R., Burkert A., 1997, MNRAS, 288, 1060
- Beech M., Mitalas R., 1994, ApJS, 95, 517
- Behrend R., Maeder A., 2001, A&A, 373, 190
- Benz W., 1990, in Numerical Modelling of Nonlinear Stellar Pulsations Problems and Prospects Smooth Particle Hydrodynamics - a Review. pp 269–+
- Bonnell I. A., Bate M. R., 2002, MNRAS, 336, 659
- Bonnell I. A., Bate M. R., Vine S. G., 2003, MNRAS, 343, 413
- Bonnell I. A., Bate M. R., Zinnecker H., 1998, MNRAS, 298, 93
- Bonnell I. A., Clarke C. J., Bate M. R., Pringle J. E., 2001, MNRAS, 324, 573
- Bonnell I. A., Vine S. G., Bate M. R., 2004, MNRAS, 349, 735
- Burkert A., Bate M. R., Bodenheimer P., 1997, MNRAS, 289, 497
- Carpenter J. M., Meyer M. R., Dougados C., Strom S. E., Hillenbrand L. A., 1997, AJ, 114, 198
- Clarke C. J., Bonnell I. A., Hillenbrand L. A., 2000, Protostars and Planets IV, pp 151–+
- Dubinski J., Narayan R., Phillips T. G., 1995, ApJ, 448, 226
- Hillenbrand L. A., Hartmann L. W., 1998, ApJ, 492, 540
- Lada C. J., Lada E. A., 2003, ARA&A, 41, 57
- Larson R. B., 1981, MNRAS, 194, 809
- Mac Low M.-M., Klessen R., Burkert A., Smith M. D., Kessel O., 1997, Bulletin of the American Astronomical Society, 29, 1244
- McKee C. F., Tan J. C., 2002, Nature, 416, 59
- McKee C. F., Tan J. C., 2003, ApJ, 585, 850
- Myers P. C., 1983, ApJ, 270, 105
- Myers P. C., Fuller G. A., 1992, ApJ, 396, 631
- Myers P. C., Gammie C. F., 1999, ApJL, 522, L141
- Norberg P., Maeder A., 2000, A&A, 359, 1025
- Padoan P., Jimenez R., Juvela M., Nordlund Å., 2004,

- ApJL, 604, L49
Shu F. H., 1977, ApJ, 214, 488
Stone J. M., Ostriker E. C., Gammie C. F., 1998, ApJL,
508, L99
Ward-Thompson D., Scott P. F., Hills R. E., Andre P.,
1994, MNRAS, 268, 276
Wolfire M. G., Cassinelli J. P., 1987, ApJ, 319, 850
Yorke H. W., Sonnhalter C., 2002, ApJ, 569, 846
Zinnecker H., McCaughrean M. J., Wilking B. A., 1993, in
Protostars and Planets III The initial stellar population.
pp 429–495

This paper has been typeset from a $\text{\LaTeX}$ file prepared
by the author.